

STAR-OPERATIONS INDUCED BY OVERRINGS.

SHARON M. CLARKE

ABSTRACT. Let D be an integral domain with quotient field K . A star-operation $\star$ on D is a closure operation $A \mapsto A^\star$ on the set of nonzero fractional ideals, $F(D)$, of D satisfying the properties: $(xD)^\star = xD$ and $(xA)^\star = xA^\star$ for all $x \in K^\star$ and $A \in F(D)$. Let $\mathcal{S}$ be a multiplicatively closed set of ideals of D . For $A \in F(D)$ define $A_{\mathcal{S}} = \{x \in K \mid xI \subseteq A, \text{ for some } I \in \mathcal{S}\}$. Then $D_{\mathcal{S}}$ is an overring of D and $A_{\mathcal{S}}$ is a fractional ideal of $D_{\mathcal{S}}$. Let $\mathcal{S}$ be a multiplicative set of finitely generated nonzero ideals of D and $A \in F(D)$, then the map $A \mapsto A_{\mathcal{S}}$ is a finite character star-operation if and only if for each $I \in \mathcal{S}$, $I_v = D$. We give an example to show that this result is not true if the ideals are not assumed to be finitely generated. In general, the map $A \mapsto A_{\mathcal{S}}$ is a star-operation if and only if $\bar{\mathcal{S}}$, the saturation of $\mathcal{S}$, is a localizing GV-system. We also discuss star-operations given of the form $A \mapsto \cap AD_{\alpha}$, where $D = \cap D_{\alpha}$.

1. INTRODUCTION

Let D be an integral domain with quotient field K . Let $K^\star = K - \{0\}$. A *fractional ideal* A of D is a D -submodule of K such that $xA \subseteq D$ for some nonzero $x \in D$. Note that the usual ideals of D , or sometimes called *integral ideals*, are also fractional ideals. Let $F(D)$ be the set of nonzero fractional ideals of D and $f(D)$ be the subset of $F(D)$ consisting of the nonzero finitely generated fractional ideals of D . Let $I(D)$ denote the set of nonzero integral ideals of D .

Definition 1.1. A **star-operation** on the set $F(D)$ is a mapping $\star : F(D) \longrightarrow F(D)$ such that for all $A, B \in F(D)$, and for all $a \in K^\star$:

- (1) $(a)^\star = (a)$ and $(aA)^\star = aA^\star$,
- (2) $A \subseteq A^\star$, and $A \subseteq B$ implies that $A^\star \subseteq B^\star$, and
- (3) $(A^\star)^\star = A^\star$.

Note that in Definition (1.1), the equality $(a)^\star = (a)$ could be replaced by the equality $D = D^\star$ since $(a)^\star = (aD)^\star = aD^\star = aD = (a)$.

For a brief introduction to star-operations, see [5, Sections 32 and 34]. However for a more detailed discussion, see [7] or [6].

For any star-operation $\star$ and $A, B \in F(D)$, we have that $(A + B)^\star = (A^\star + B^\star)^\star$ and $(AB)^\star = (AB^\star)^\star = (A^\star B)^\star$. A star-operation $\star$ *distributes over intersections* if $(A \cap B)^\star = A^\star \cap B^\star$ for all $A, B \in F(D)$. This is easily seen to be equivalent to $(A_1 \cap \dots \cap A_n)^\star = A_1^\star \cap \dots \cap A_n^\star$ for all (integral) $A_1, \dots, A_n \in F(D)$. We say that a star-operation $\star$ *distributes over arbitrary intersections* if $0 \neq \cap A_\alpha \Rightarrow (\cap A_\alpha)^\star = \cap A_\alpha^\star$ for all $A \in F(D)$.

A star-operation $\star$ has *finite character* if, for each $A \in F(D)$, we have that $A^\star = \cup\{B^\star \mid B \in f(D) \text{ and } B \subseteq A\}$. If $\star$ is a star-operation on $F(D)$, we can define a finite character star-operation $\star_s$ by $A \mapsto A^{\star_s} = \cup\{B^\star \mid B \in f(D) \text{ and } B \subseteq A\}$. If $A \in f(D)$, then $A^\star = A^{\star_s}$. Three important examples of star-operations are the *d-operation*, the *v-operation*, and the *t-operation*. The *d-operation* is the identity operation $A_d = A$ for all $A \in F(D)$. The *v-operation* is defined by $A \mapsto A_v = (A^{-1})^{-1}$, where $A^{-1} = [D : A] = \{x \in K \mid xA \subseteq D\}$, or equivalently $A \mapsto A_v = \cap\{(x) \mid A \subseteq (x), 0 \neq x \in K\}$. Throughout this paper, for $A, B \in F(D)$, $[A : B] = \{x \in K \mid xB \subseteq A\}$, and $(A : B) = \{x \in D \mid xB \subseteq A\} = [A : B] \cap D$. Note that for any star-operation $\star$, $A^\star \subseteq A_v$ for all $A \in F(D)$. The *t-operation* is defined by $A \mapsto A_t = \cup\{B_v \mid B \in f(D) \text{ and } B \subseteq A\}$. Note that $t = v_s$. Other examples of star-operations will be given and defined in this paper.

Suppose that $\star_1$ and $\star_2$ are two star-operations on $F(D)$, then $\star_1 \leq \star_2$ if $A^{\star_1} \subseteq A^{\star_2}$ for all $A \in F(D)$ (or equivalently, $F_{\star_2}(D) \subseteq F_{\star_1}(D)$), and $\star_1 = \star_2$ if $A^{\star_1} = A^{\star_2}$ for all $A \in F(D)$. The star-operations $\star_1$ and $\star_2$ are *equivalent* if $A^{\star_1} = A^{\star_2}$ for all $A \in f(D)$. According to [5], for any star-operation $\star$, $d \leq \star \leq v$.

A ring D' is said to be an *overring* of D if $D \subseteq D' \subseteq K$. Let $\mathcal{S}$ be a multiplicatively closed set of ideals (or a multiplicative set of ideals) of D , i.e, $I, J \in \mathcal{S}$ implies that $IJ \in \mathcal{S}$. For a fractional ideal A of D , define $A_{\mathcal{S}} = \{x \in K \mid xI \subseteq A, \text{ for some } I \in \mathcal{S}\}$. Then $D_{\mathcal{S}}$ is an overring of D , called the *$\mathcal{S}$ -transform* of D , and $A_{\mathcal{S}}$ is a fractional ideal of $D_{\mathcal{S}}$. In [4], Arnold and Brewer discuss the relationship between the ideal structure of a commutative ring R and that of the $\mathcal{S}$ -transform of R .

Let $\mathcal{S}$ be a multiplicative set of ideals of D . The *saturation*, $\bar{\mathcal{S}}$, of $\mathcal{S}$ is defined as $\bar{\mathcal{S}} = \{A \mid I \subseteq A \subseteq D, \text{ for some } I \in \mathcal{S}\}$. We say that $\mathcal{S}$ is saturated if $\mathcal{S} = \bar{\mathcal{S}}$. Note that $\bar{\bar{\mathcal{S}}} = \bar{\mathcal{S}}$ so the saturation of a multiplicative set of ideals is saturated. It is easily proved that $I, J \in \bar{\mathcal{S}}$ implies that $IJ \in \bar{\mathcal{S}}$. So $\bar{\mathcal{S}}$ is a multiplicative set of ideals. Clearly $A_{\mathcal{S}} = A_{\bar{\mathcal{S}}}$.

Definition 1.2. A nonempty set $\mathcal{F}$ of ideals of D is a **localizing system** if

- (1) $I \in \mathcal{F}$ and J an ideal with $I \subseteq J \subseteq D$ imply that $J \in \mathcal{F}$, and
- (2) $I \in \mathcal{F}$ and J an ideal with $(J : i) \in \mathcal{F}$ for all $i \in I$ imply that $J \in \mathcal{F}$.

In Section 2, we investigate when the map $A \mapsto A_{\mathcal{S}}$ is a star-operation. This question is answered for $\mathcal{S}$ a multiplicative set of ideals of D and for $\mathcal{S}$ a multiplicative set of finitely generated ideals of D . An obvious necessary condition is that $D_{\mathcal{S}} = D$. We also determine when this star-operation has finite character and we give an example where the map $A \mapsto A_{\mathcal{S}}$ is not a star-operation.

In Section 3, we characterize when star-operations of the form $A \mapsto \cap AD_{\alpha}$, with $D = \cap D_{\alpha}$, have finite character.

I give an idea of what I will be working on in the near future in Section 4.

2. STAR-OPERATIONS AND GENERALIZED QUOTIENT RINGS

Recall that a star-operation has finite character if for each $A \in F(D)$, $A^{\star} = \cup\{B^{\star} \mid B \in f(D) \text{ and } B \subseteq A\}$.

Note that the d -operation and the t -operation (defined in the introduction) have finite character, but the v -operation need not have finite character.

Definition 2.1. Let $\mathcal{S}$ be a multiplicatively closed set of ideals of D . Then for $A \in F(D)$, $A_{\mathcal{S}} = \{x \in K \mid xI \subseteq A, \text{ for some } I \in \mathcal{S}\}$.

Recall also that $D_{\mathcal{S}}$ is an overring of D , and if A is a fractional ideal of D , then $A_{\mathcal{S}}$ is a fractional ideal of $D_{\mathcal{S}}$.

Definition 2.2. Let D be an integral domain and $\star$ a star-operation on $F(D)$. Then we define $\mathcal{S}_{\star} = \{I \in I(D) \mid I^{\star} = D\}$ and $\mathcal{S}_{\star}^f = \{I \in \mathcal{S}_{\star} \mid I \text{ is finitely generated}\}$. In particular, $\mathcal{S}_v = \{I \in I(D) \mid I_v = D \text{ or } I^{-1} = D\}$ and $\mathcal{S}_v^f = \{I \in \mathcal{S}_v \mid I \text{ is finitely generated}\}$.

Definition 2.3. Let $I \in F(D)$, then I is said to be a **GV-ideal** (Glaz-Vasconcelos ideal) if $I_v = D$, or equivalently $I^{-1} = D$. A multiplicative set $\mathcal{S}$ is said to be a **GV-system** if $\mathcal{S} \subseteq \mathcal{S}_v$.

Proposition 2.4. $D = D_{\mathcal{S}}$ if and only if $\mathcal{S} \subseteq \mathcal{S}_v$.

Proof. ($\Rightarrow$) Let $I \in \mathcal{S}$. Since $I^{-1}I \subseteq D$, $I^{-1} \subseteq D_{\mathcal{S}} = D$. We know that $D \subseteq I^{-1}$. Therefore $I^{-1} = D$. So $I \in \mathcal{S}_v$. Hence $\mathcal{S} \subseteq \mathcal{S}_v$.

($\Leftarrow$) Now $D \subseteq D_{\mathcal{S}}$. Let $x \in D_{\mathcal{S}}$, then there exists $I \in \mathcal{S} \subseteq \mathcal{S}_v$ such that $xI \subseteq D$. This implies that $x \in [D : I] = I^{-1} = D$ since $I \in \mathcal{S}_v$. Therefore $D_{\mathcal{S}} \subseteq D$, and hence $D = D_{\mathcal{S}}$. $\square$

Proposition 2.5. *Let $\mathcal{S}$ be a multiplicative set of nonzero ideals of D . Let $A, B, A_1, \dots, A_n \in F(D)$ and $a \in K^*$, then*

- (1) $A \subseteq B$ implies that $A_{\mathcal{S}} \subseteq B_{\mathcal{S}}$,
- (2) $A \subseteq A_{\mathcal{S}}$,
- (3) $(aA)_{\mathcal{S}} = aA_{\mathcal{S}}$,
- (4) $(A_1 \cap \dots \cap A_n)_{\mathcal{S}} = A_{1\mathcal{S}} \cap \dots \cap A_{n\mathcal{S}}$,
- (5) If A and B are integral ideals of D with $A_{\mathcal{S}} = B_{\mathcal{S}} = D_{\mathcal{S}}$, then $(AB)_{\mathcal{S}} = D_{\mathcal{S}}$, and
- (6) The following are equivalent:
 - (a) $\mathcal{S} \subseteq \mathcal{S}_v$,
 - (b) $D = D_{\mathcal{S}}$, and
 - (c) $(a)_{\mathcal{S}} = (a)$ for all $a \in K^*$.

Proof. (1) Let $x \in A_{\mathcal{S}}$, then there exists $I \in \mathcal{S}$ such that $xI \subseteq A \subseteq B$. Therefore $x \in B_{\mathcal{S}}$ by definition. Hence $A_{\mathcal{S}} \subseteq B_{\mathcal{S}}$.

(2) Let $x \in A$ and $I \in \mathcal{S}$, then $xI \subseteq AD \subseteq A$. So $x \in A_{\mathcal{S}}$, and therefore $A \subseteq A_{\mathcal{S}}$.

(3) Let $x \in (aA)_{\mathcal{S}}$, then there exists $I \in \mathcal{S}$ such that $xI \subseteq aA$. Since K is a field and $a \in K^*$, then $\frac{x}{a}I \subseteq A$. So $\frac{x}{a} \in A_{\mathcal{S}}$. Therefore $x \in aA_{\mathcal{S}}$. So $(aA)_{\mathcal{S}} \subseteq aA_{\mathcal{S}}$.

Suppose $x \in aA_{\mathcal{S}}$, then $\frac{x}{a} \in A_{\mathcal{S}}$ since K is a field and $a \in K^*$. Therefore there exists $I \in \mathcal{S}$ such that $\frac{x}{a}I \subseteq A$. This implies that $xI \subseteq aA$ so $x \in (aA)_{\mathcal{S}}$. Therefore $aA_{\mathcal{S}} \subseteq (aA)_{\mathcal{S}}$, and hence $(aA)_{\mathcal{S}} = aA_{\mathcal{S}}$.

(4) Since $A_1 \cap \dots \cap A_n \subseteq A_i$ for each $i = 1, \dots, n$, then by (1) $(A_1 \cap \dots \cap A_n)_{\mathcal{S}} \subseteq A_{i\mathcal{S}}$ for each $i = 1, \dots, n$. Therefore $(A_1 \cap \dots \cap A_n)_{\mathcal{S}} \subseteq A_{1\mathcal{S}} \cap \dots \cap A_{n\mathcal{S}}$.

Let $x \in A_{1\mathcal{S}} \cap \dots \cap A_{n\mathcal{S}}$, then $x \in A_{i\mathcal{S}}$ for each $i = 1, \dots, n$. Therefore there exist $I_1, \dots, I_n \in \mathcal{S}$ such that $xI_1 \subseteq A_1, \dots, xI_n \subseteq A_n$. So for each A_i we have $xI_1 \cdots I_n = xI_1 \cdots I_i \cdots I_n \subseteq xI_i \subseteq A_i$. Therefore $xI_1 \cdots I_n \subseteq A_1 \cap \dots \cap A_n$. Since $\mathcal{S}$ is multiplicatively closed, $I_1 \cdots I_n \in \mathcal{S}$. Therefore $x \in (A_1 \cap \dots \cap A_n)_{\mathcal{S}}$. Thus $A_{1\mathcal{S}} \cap \dots \cap A_{n\mathcal{S}} \subseteq (A_1 \cap \dots \cap A_n)_{\mathcal{S}}$, and hence $(A_1 \cap \dots \cap A_n)_{\mathcal{S}} = A_{1\mathcal{S}} \cap \dots \cap A_{n\mathcal{S}}$.

(5) Suppose that $A, B \in I(D)$. Then $AB \subseteq D$. Therefore $(AB)_{\mathcal{S}} \subseteq D_{\mathcal{S}}$.

Since $A_{\mathcal{S}} = D_{\mathcal{S}}$, $1 \in A_{\mathcal{S}}$. This implies that there exists $I \in \mathcal{S}$ such that $1I \subseteq A$. Similarly, $B_{\mathcal{S}} = D_{\mathcal{S}}$ implies that there exists $J \in \mathcal{S}$ such that $1J \subseteq B$. So $1IJ = 1I1J \subseteq AB$. Therefore since $IJ \in \mathcal{S}$, $1 \in (AB)_{\mathcal{S}}$. So $D_{\mathcal{S}} \subseteq (AB)_{\mathcal{S}}$. Hence $(AB)_{\mathcal{S}} = D_{\mathcal{S}}$.

(6) (a) $\Leftrightarrow$ (b) This is true by Proposition 2.4.

(b) $\Rightarrow$ (c) Suppose $D = D_{\mathcal{S}}$, and let $a \in K^*$. Then $(a) = aD = aD_{\mathcal{S}} = (aD)_{\mathcal{S}} = (a)_{\mathcal{S}}$.

(c) $\Rightarrow$ (b) $1 \in K^*$. So take $a = 1$. Then by (c), $(1)_{\mathcal{S}} = (1)$. Therefore $D_{\mathcal{S}} = (1)_{\mathcal{S}} = (1) = D$. $\square$

We now state our first theorem which characterizes when the map $A \mapsto A_{\mathcal{S}}$, for $A \in F(D)$ and $\mathcal{S}$ a multiplicative set of finitely generated nonzero ideals, is a star-operation.

Theorem 2.6. *Let $\mathcal{S}$ be a multiplicative set of finitely generated nonzero ideals of D . The map $A \mapsto A_{\mathcal{S}}$ ($A \in F(D)$) is a star-operation if and only if $\mathcal{S} \subseteq \mathcal{S}_v$. In this case the star-operation $A \mapsto A_{\mathcal{S}}$ has finite character.*

Proof. ($\Rightarrow$) Suppose the map $A \mapsto A_{\mathcal{S}}$ is a star-operation, then $D = D_{\mathcal{S}}$. Therefore $\mathcal{S} \subseteq \mathcal{S}_v$, by Proposition 2.5.

($\Leftarrow$) Since $\mathcal{S}$ is a multiplicative set of nonzero ideals, then by Proposition (2.5), $A \subseteq A_{\mathcal{S}}$ (and thus $A_{\mathcal{S}} \subseteq (A_{\mathcal{S}})_{\mathcal{S}}$), $(xA)_{\mathcal{S}} = xA_{\mathcal{S}}$, and $A \subseteq B$ implies that $A_{\mathcal{S}} \subseteq B_{\mathcal{S}}$. Also by Proposition 2.5, since $\mathcal{S} \subseteq \mathcal{S}_v$, then $(x)_{\mathcal{S}} = (x)$ for all $x \in K^*$. So it suffices to show that $(A_{\mathcal{S}})_{\mathcal{S}} \subseteq A_{\mathcal{S}}$.

Let $x \in (A_{\mathcal{S}})_{\mathcal{S}}$, then there exists $I \in \mathcal{S}$ such that $xI \subseteq A_{\mathcal{S}}$. I is finitely generated so suppose that $I = (x_1, \dots, x_n)$. Then for each $x_i \in I$, $xx_i \in A_{\mathcal{S}}$. Therefore for each x_i , there exists $A_i \in \mathcal{S}$ such that $xx_i A_i \subseteq A$. This implies that $xx_i \prod_{i=1}^n A_i \subseteq xx_i A_i \subseteq A$ for each x_i . So $xI \prod_{i=1}^n A_i \subseteq A$ for each i . Now $I \prod_{i=1}^n A_i \in \mathcal{S}$ since $\mathcal{S}$ is multiplicatively closed. Therefore $x \in A_{\mathcal{S}}$, and hence $(A_{\mathcal{S}})_{\mathcal{S}} \subseteq A_{\mathcal{S}}$. So the map $A \mapsto A_{\mathcal{S}}$ is a star-operation.

We now proceed to show that the map $A \mapsto A_{\mathcal{S}}$ has finite character (i.e., show that $A_{\mathcal{S}} = \cup\{B_{\mathcal{S}} \mid B \in f(D) \text{ and } B \subseteq A\}$).

Let $x \in \cup\{B_{\mathcal{S}} \mid B \in f(D) \text{ and } B \subseteq A\}$, then $x \in B_{\mathcal{S}}$ such that $B \in f(D)$ and $B \subseteq A$ for some B . But $B \subseteq A$ implies that $B_{\mathcal{S}} \subseteq A_{\mathcal{S}}$. Therefore $x \in A_{\mathcal{S}}$, and hence $\cup\{B_{\mathcal{S}} \mid B \in f(D) \text{ and } B \subseteq A\} \subseteq A_{\mathcal{S}}$.

Let $x \in A_{\mathcal{S}}$, then there exists $I \in \mathcal{S}$ such that $xI \subseteq A$. Since I is finitely generated, xI is a finitely generated fractional ideal contained in A . So if we let $B = xI$, then $xI = B$ implies that $x \in B_{\mathcal{S}}$. So $x \in \cup\{B_{\mathcal{S}} \mid B \in f(D) \text{ and } B \subseteq A\}$. Therefore $A_{\mathcal{S}} \subseteq \cup\{B_{\mathcal{S}} \mid B \in f(D), B \subseteq A\}$, and hence $A_{\mathcal{S}} = \cup\{B_{\mathcal{S}} \mid B \in f(D) \text{ and } B \subseteq A\}$. Thus the map $A \mapsto A_{\mathcal{S}}$ has finite character. $\square$

Note that the above result is false if the ideals in $\mathcal{S}$ are not assumed to be finitely generated as the following example will show. But first, we state a corollary to Theorem 2.6

Corollary 2.7. *Let $\mathcal{S}$ be a multiplicative set of finitely generated nonzero GV-ideals of D . Then the map $A \mapsto A_{\mathcal{S}} = A^*$ is a finite character star-operation on D that distributes over finite intersections.*

Proof. Since $\mathcal{S}$ is a multiplicative set of GV-ideals, $\mathcal{S} \subseteq \mathcal{S}_v$ by definition. Also $\mathcal{S}$ consists of finitely generated ideals by hypothesis. Therefore by Theorem 2.6, the map $A \mapsto A_{\mathcal{S}} = A^*$ is a finite character star-operation on D . By Proposition 2.5 we have that the map $A \mapsto A_{\mathcal{S}} = A^*$ distributes over finite intersections. $\square$

Example 2.8. Let $D = K[\{x_n\}_{n=1}^{\infty}]$, where K is a field, and let $M = (\{x_n\}_{n=1}^{\infty})$. Then M is a maximal ideal of D . Now $M_v = D$. Let $\mathcal{S} = \{M^n\}_{n=1}^{\infty}$. Now $M_v = D$ implies that $(M^n)_v = D$ for each n . So $\mathcal{S} \subseteq \mathcal{S}_v$. Therefore $D = D_{\mathcal{S}}$ by Proposition 2.5. Let $A = (\{x_n M^n\}_{n=1}^{\infty})$. Now $x_n M^n \subseteq A$ so $x_n \in A_{\mathcal{S}}$. Hence $M = (\{x_n\}_{n=1}^{\infty}) \subseteq A_{\mathcal{S}}$. Now $1 \notin A_{\mathcal{S}}$ for $1M^n = M^n \not\subseteq A$ for any n . So since $M \subseteq A_{\mathcal{S}} \subsetneq D$ and M is a maximal ideal of D , $M = A_{\mathcal{S}}$. Therefore $M_{\mathcal{S}} = (A_{\mathcal{S}})_{\mathcal{S}}$.

Now $M \in \mathcal{S}$, and $1M \subseteq M$. Therefore $1 \in M_{\mathcal{S}}$ and thus $M_{\mathcal{S}} = D_{\mathcal{S}}$. So $(A_{\mathcal{S}})_{\mathcal{S}} = M_{\mathcal{S}} = D_{\mathcal{S}} = D$. Hence $A_{\mathcal{S}} = M \subsetneq D = (A_{\mathcal{S}})_{\mathcal{S}}$. Therefore in this example, the map $A \mapsto A_{\mathcal{S}}$ is not a star-operation.

Next, we give a more general characterization of Theorem 2.6. But first, we look at some useful results.

Recall that for, $\mathcal{S}$, a multiplicative set of ideals of D , the saturation of $\mathcal{S}$ is defined by $\bar{\mathcal{S}} = \{A \mid I \subseteq A \subseteq D, \text{ where } I \in \mathcal{S}\}$. Clearly $\mathcal{S} \subseteq \bar{\mathcal{S}}$ since for any $I \in \mathcal{S}$, $I \subseteq I \subseteq D$, which implies that $I \in \bar{\mathcal{S}}$.

Definition 2.9. A nonempty set $\mathcal{F}$ of ideals of D is a **localizing system** if

- (1) $I \in \mathcal{F}$ and J an ideal with $I \subseteq J \subseteq D$ imply that $J \in \mathcal{F}$, and
- (2) $I \in \mathcal{F}$ and J an ideal with $(J : i) \in \mathcal{F}$ for all $i \in I$ imply that $J \in \mathcal{F}$.

Proposition 2.10. Let $\mathcal{S}$ be a multiplicative set of ideals of D . Then for $I \in I(D)$, $I_{\mathcal{S}} = D_{\mathcal{S}}$ if and only if $I \in \bar{\mathcal{S}}$.

Proof. $(\Rightarrow)$ $I_{\mathcal{S}} = D_{\mathcal{S}}$ implies that $1 \in I_{\mathcal{S}}$. So there exists $J \in \mathcal{S}$ such that $1J \subseteq I$. Therefore $J = 1J \subseteq I \subseteq D$. Thus $I \in \bar{\mathcal{S}}$.

$(\Leftarrow)$ Suppose that $I \in \bar{\mathcal{S}}$. Then there exists $J \in \mathcal{S}$ such that $J \subseteq I \subseteq D$. This implies that $1J = J \subseteq I$. Therefore $1 \in I_{\mathcal{S}}$. So $D_{\mathcal{S}} \subseteq I_{\mathcal{S}}$. Hence $I_{\mathcal{S}} = D_{\mathcal{S}}$. $\square$

Theorem 2.11. Let $\mathcal{S}$ be a multiplicative set of ideals of D . Then the following are equivalent:

- (1) $A_{\mathcal{S}} = (A_{\mathcal{S}})_{\mathcal{S}}$ for each $A \in F(D)$,
- (2) $I_{\mathcal{S}} = (I_{\mathcal{S}})_{\mathcal{S}}$ for each $I \in I(D)$, and
- (3) $\mathcal{S}$ is a localizing system.

Proof. (1) $\Rightarrow$ (2) This is clear since $I(D) \subseteq F(D)$.

(2) $\Rightarrow$ (3) Suppose that $I \in \bar{\mathcal{S}}$ and $I \subseteq J \subseteq D$ for some ideal J in D . Then by definition, $J \in \bar{\bar{\mathcal{S}}} = \bar{\mathcal{S}}$.

Now suppose that $I \in \bar{\mathcal{S}}$ and $J \in I(D)$ with $(J : i) \in \bar{\mathcal{S}}$ for all $i \in I$. Thus $(J : i)_{\mathcal{S}} = D_{\mathcal{S}}$ by Proposition 2.10. So $(J : i)i \subseteq J$ gives $(J : i)_{\mathcal{S}}(i)_{\mathcal{S}} \subseteq J_{\mathcal{S}}$ and hence for each $i \in I$, $i \in (i)_{\mathcal{S}} = (i)_{\mathcal{S}}D_{\mathcal{S}} = (J : i)_{\mathcal{S}}(i)_{\mathcal{S}} \subseteq J_{\mathcal{S}}$. This implies that $I \subseteq J_{\mathcal{S}}$. Thus $I_{\mathcal{S}} \subseteq (J_{\mathcal{S}})_{\mathcal{S}} = J_{\mathcal{S}}$. The last equality follows from the hypothesis. Now by Proposition 2.10, $I_{\mathcal{S}} = D_{\mathcal{S}}$ so $D_{\mathcal{S}} = I_{\mathcal{S}} \subseteq J_{\mathcal{S}} \subseteq D_{\mathcal{S}}$. Hence $J_{\mathcal{S}} = D_{\mathcal{S}}$, and so again by Proposition 2.10, $J \in \bar{\mathcal{S}}$. So $\bar{\mathcal{S}}$ is a localizing system.

(3) $\Rightarrow$ (1) Suppose that $\bar{\mathcal{S}}$ is a localizing system and let $A \in F(D)$. We know that $A_{\mathcal{S}} \subseteq (A_{\mathcal{S}})_{\mathcal{S}}$. Let $x \in (A_{\mathcal{S}})_{\mathcal{S}}$. Then there exists $I \in \mathcal{S}$ such that $xI \subseteq A_{\mathcal{S}}$. Let $J = (A : x)$. Then $J \in I(D)$. Now for $i \in I$, $xi \in A_{\mathcal{S}}$. So there exists $J_i \in \mathcal{S}$ such that $xiJ_i \subseteq A$. So $iJ_i \subseteq (A : x) = J$, that is $J_i \subseteq (J : i)$. Hence $(J : i) \in \bar{\mathcal{S}}$ for each $i \in I$ since $\bar{\mathcal{S}}$ is a localizing system. So $J \in \bar{\mathcal{S}}$, again since $\bar{\mathcal{S}}$ is a localizing system. Now $xJ \subseteq A$ so $x \in A_{\bar{\mathcal{S}}} = A_{\mathcal{S}}$. Thus $(A_{\mathcal{S}})_{\mathcal{S}} \subseteq A_{\mathcal{S}}$. Hence $A_{\mathcal{S}} = (A_{\mathcal{S}})_{\mathcal{S}}$. $\square$

We now state the more general characterization of Theorem 2.6

Theorem 2.12. *Let $\mathcal{S}$ be a multiplicative set of ideals. Then for $A \in F(D)$, the map $A \mapsto A_{\mathcal{S}}$ is a star-operation if and only if $\bar{\mathcal{S}}$ is a localizing GV-system.*

Proof. ($\Rightarrow$) Suppose that the map $A \mapsto A_{\mathcal{S}}$ is a star-operation for $A \in F(D)$. Then $A_{\mathcal{S}} = (A_{\mathcal{S}})_{\mathcal{S}}$. Therefore $\bar{\mathcal{S}}$ is a localizing system by Theorem 2.11.

Let $I \in \bar{\mathcal{S}}$. Then there exists $J \in \mathcal{S}$ such that $J \subseteq I \subseteq D$. Since the map $A \mapsto A_{\mathcal{S}}$ is a star-operation, $(x)_{\mathcal{S}} = (x)$ for all $x \in K^*$. So $\mathcal{S} \subseteq \mathcal{S}_v$ by Proposition 2.5. This implies that $J_v = D$. So $D = J_v \subseteq I_v \subseteq D_v = D$. Therefore $I_v = D$ which implies that $I \in \mathcal{S}_v$. Thus $\bar{\mathcal{S}} \subseteq \mathcal{S}_v$. Hence $\bar{\mathcal{S}}$ is a localizing GV-system.

($\Leftarrow$) Since $\bar{\mathcal{S}}$ is a GV-system, $\bar{\mathcal{S}} \subseteq \mathcal{S}_v$. So $\mathcal{S} \subseteq \bar{\mathcal{S}} \subseteq \mathcal{S}_v$, and thus $D = D_{\mathcal{S}}$ by Proposition 2.5.

Since $\bar{\mathcal{S}}$ is a localizing system, $A_{\mathcal{S}} = (A_{\mathcal{S}})_{\mathcal{S}}$ for all $A \in F(D)$ by Theorem 2.11. The other conditions for a star-operation are satisfied by Proposition 2.5. Hence the map $A \mapsto A_{\mathcal{S}}$ is a star-operation. $\square$

3. STAR-OPERATIONS INDUCED BY THE LOCALIZATIONS OF D

In this section, we study the star-operations that are induced by localizations $D_{P_{\alpha}}$ of D at P_{α} where $\{P_{\alpha}\}$ is a collection of nonzero prime ideals of D . We then extend our study to the case where the star-operations are induced by arbitrary overrings of D .

Lemma 3.1. *Let $A, B \in F(D)$. Then $(A : B)B \subseteq A \cap B$. If B is principal, then $(A : B)B = A \cap B$.*

Proof. Let $a \in (A : B)$. Then $aB \subseteq A$. Therefore $(A : B)B \subseteq A$. Suppose $a \in (A : B)$ and $b \in B$. Then $ab \in B$ since $a \in D$. Therefore $(A : B)B \subseteq B$. Thus $(A : B)B \subseteq A \cap B$.

Suppose B is principal, then $B = (x)$ for some $x \in K^*$. Let $a \in A \cap B$. Then $a \in A$ and $a \in B$. Now $a \in B = (x)$ implies that $a = bx$ for some $b \in D$, and $a \in A$ implies that $bx = a \in A$. Therefore $b(x) = b(xD) = (bx)D \subseteq AD \subseteq A$. So $b \in (A : (x))$. This implies that $a = bx \in (A : (x))(x) = (A : B)B$, and thus $A \cap B \subseteq (A : B)B$. Hence if B is principal, then $(A : B)B = A \cap B$. $\square$

Theorem 3.2. *Let $A, B \in F(D)$, $x \in K^*$ and $\star : F(D) \longrightarrow F(D)$ an operation satisfying:*

- (1) $A \subseteq B$ implies $A^\star \subseteq B^\star$,
- (2) $(xA)^\star = xA^\star$,
- (3) $(A \cap B)^\star = A^\star \cap B^\star$,
- (4) $A^\star = B^\star = D$ implies $(AB)^\star = D$, and
- (5) $D^\star = D$ (or equivalently, $(x)^\star = (x)$ for all $x \in K^*$).

Then $\mathcal{S}_\star = \{I \in I(D) \mid I^\star = D\}$ is a multiplicative set of nonzero ideals of D , and $A^\star = A_{\mathcal{S}_\star}$.

Proof. Let $I, J \in \mathcal{S}_\star$. Then $I^\star = D = J^\star$. Therefore $(IJ)^\star = D$ by (4). So $IJ \in \mathcal{S}_\star$, and thus $\mathcal{S}_\star$ is a multiplicative set of nonzero ideals of D .

Let $x \in A_{\mathcal{S}_\star}$. Then there exists $I \in \mathcal{S}_\star$ such that $xI \subseteq A$. Since $I \in \mathcal{S}_\star$, $I^\star = D$. Therefore $x \in xD = xI^\star = (xI)^\star \subseteq A^\star$. The second equality follows from (2), and the last containment follows from (1). So $A_{\mathcal{S}_\star} \subseteq A^\star$.

Let $x \in A^\star$, then $(x) \subseteq A^\star$. So $(x) = A^\star \cap (x) = A^\star \cap (x)^\star = (A \cap (x))^\star = [(A : (x))(x)]^\star = (A : (x))^\star(x)$. The second and third equalities are due to (5) and (3) respectively, and the fourth equality is due to Lemma 3.1. So $(A : (x))^\star = D$, and thus $(A : (x)) \in \mathcal{S}_\star$. Now $x(A : (x)) = (x)(A : (x)) = A \cap (x) \subseteq A$. So $x \in A_{\mathcal{S}_\star}$, and therefore $A^\star \subseteq A_{\mathcal{S}_\star}$. Hence $A^\star = A_{\mathcal{S}_\star}$. $\square$

We now state our first major theorem of this section.

Theorem 3.3. *Let $D = \cap D_{P_\alpha}$ where $\{P_\alpha\}$ is a set of nonzero prime ideals of D . Let $\star$ be the star-operation defined by $A^\star = \cap AD_{P_\alpha}$. Then the following are equivalent:*

- (1) *The map $A \longmapsto \cap AD_{P_\alpha} = A^\star$ has finite character,*
- (2) *$A^\star = D$ implies that there exists a finitely generated ideal $B \subseteq A$ with*

$B^* = D$, and

(3) If A is an ideal of D with $A \not\subseteq P_\alpha$ for each α , then there exists a finitely

generated ideal $B \subseteq A$ with $B \not\subseteq P_\alpha$ for each α .

Proof. (1) $\Rightarrow$ (2) Let $A \in I(D)$. Since $\star$ has finite character, $A^* = \cup\{B^* \mid B \in f(D) \text{ and } B \subseteq A\}$. So if $A^* = D$, then $1 \in D = A^* = \cup\{B^* \mid B \in f(D) \text{ and } B \subseteq A\}$. So $1 \in B^*$ for some B^* where B is finitely generated. Therefore $B^* = D$ for some finitely generated ideal $B \subseteq A$. (The proof shows that this holds for any finite character star-operation.)

(2) $\Rightarrow$ (1) Now $\star$ distributes over finite intersections. Therefore by Theorem 3.2, for $\mathcal{S}_\star$ a multiplicative subset of D , $A^* = A_{\mathcal{S}_\star}$ for all $A \in F(D)$, and $\mathcal{S}_\star \subseteq \mathcal{S}_v$.

Claim 1. $A_{\mathcal{S}_\star} = A_{\mathcal{S}_\star^f}$ for all $A \in F(D)$.

We already know that $\mathcal{S}_\star^f \subseteq \mathcal{S}_\star$. So let $A \in F(D)$. Then $A_{\mathcal{S}_\star^f} \subseteq A_{\mathcal{S}_\star}$. Let $x \in A_{\mathcal{S}_\star}$. Then $xI \subseteq A$ for some $I \in I(D)$ such that $I^* = D$. So by (2), there exists a finitely generated ideal $J \subseteq I$ with $J^* = D$. Thus $xJ \subseteq xI \subseteq A$. But $J \in \mathcal{S}_\star^f$. Therefore $x \in A_{\mathcal{S}_\star^f}$. This implies that $A_{\mathcal{S}_\star} \subseteq A_{\mathcal{S}_\star^f}$. Hence $A_{\mathcal{S}_\star} = A_{\mathcal{S}_\star^f}$.

Now $A_{\mathcal{S}_\star} = A_{\mathcal{S}_\star^f}$ for all $A \in F(D)$. Therefore for all $A \in F(D)$, $A^* = A_{\mathcal{S}_\star} = A_{\mathcal{S}_\star^f}$ by Claim 1, and $\mathcal{S}_\star^f \subseteq \mathcal{S}_\star \subseteq \mathcal{S}_v$. So by Corollary 2.7, the map $A \mapsto \cap AD_{P_\alpha} = A^*$ has finite character.

(2) $\Rightarrow$ (3) Let $A \in I(D)$ such that $A \not\subseteq P_\alpha$ for each α . Then $A^* = D$ by [1, Theorem 1]. Therefore by (2), there exists a finitely generated ideal $B \subseteq A$ such that $B^* = D$. But by [1, Theorem 1], $P_\alpha^* = P_\alpha$ for each $P_\alpha \in \{P_\alpha\}$. So for each P_α , $P_\alpha^* = P_\alpha \subsetneq D = B^*$. Suppose that $B \subseteq P_\alpha$ for some α , then $B^* \subseteq P_\alpha^*$ which is a contradiction. Therefore $B \not\subseteq P_\alpha$ for each α .

(3) $\Rightarrow$ (2) Suppose that $A \in I(D)$ with $A^* = D$. Now $P_\alpha^* = P_\alpha \subsetneq D = A^*$ for each $P_\alpha \in \{P_\alpha\}$. The first equality follows from [1, Theorem 1]. Suppose that $A \subseteq P_\alpha$ for some α . Then $A^* \subseteq P_\alpha^*$ which is again a contradiction. So $A \not\subseteq P_\alpha$ for each α . Therefore by (3), there exists a finitely generated ideal $B \subseteq A$ with $B \not\subseteq P_\alpha$ for each α . But according to [1, Theorem 1], this implies that $B^* = D$. $\square$

I am now considering under what conditions Theorem 3.3 can be extended to star-operations of the form $A \mapsto \cap AD_\alpha$ where $D = \cap D_\alpha$. I have the following result.

Let D_α be an overring of D . According to [1], a star-operation of the form $A \mapsto \cap AD_\alpha$, where $D = \cap D_\alpha$, is just the special case of the

star-operation, $\star_\alpha$, defined by the map $A \mapsto A^\star = \cap(AD_\alpha)^{\star_\alpha}$, where each $\star_\alpha$ (the star-operation on the corresponding D_α) is the identity operation d on D_α .

Definition 3.4. A right R -module M is **flat** if and only if for every exact sequence of R -modules,

$$0 \longrightarrow N' \xrightarrow{f} N \xrightarrow{g} N'' \longrightarrow 0,$$

the sequence

$$0 \longrightarrow M \otimes_R N' \xrightarrow{1_M \otimes f} M \otimes_R N \xrightarrow{1_M \otimes g} M \otimes_R N'' \longrightarrow 0$$

is exact.

Definition 3.5. An overring T of a ring R is a **flat overring** if T is a flat R -module.

According to [8, Theorem 3.3], a localization is always flat.

Theorem 3.6. Let $D = \cap D_\alpha$ where each D_α is a flat overring of D . Then for $A \in F(D)$, the following are equivalent:

- (1) The map $A \mapsto \cap AD_\alpha = A^\star$ has finite character, and
- (2) $A^\star = D$ implies that there exists a finitely generated ideal $B \subseteq A$ with

$$B^\star = D.$$

Proof. (1) $\Rightarrow$ (2) This follows from Theorem 3.3.

(2) $\Rightarrow$ (1) Suppose that $A^\star = D$ where $A^\star = \cap AD_\alpha$. Recall that the operation $A \mapsto A^\star = \cap AD_\alpha$ is a special case of the star-operation $\star_\alpha$, defined earlier, where each $\star_\alpha$ is the identity operation d on D_α . So let $A, B \in F(D_\alpha)$. Then $(A \cap B)^{\star_\alpha} = (A \cap B)_d = A \cap B = A_d \cap B_d = A^{\star_\alpha} \cap B^{\star_\alpha}$. Since each D_α is flat, $(A \cap B)^\star = A^\star \cap B^\star$ for all $A, B \in F(D)$ by [1, Theorem 2]. That is, $\star$ distributes over intersections. We now proceed in the same way as in the proof of Theorem 3.3 ((2) $\Rightarrow$ (1)) to conclude that the map $A \mapsto \cap AD_\alpha = A^\star$ has finite character. $\square$

4. FUTURE WORK

In his paper, “Star-Operations Induced by Overrings”, D. D. Anderson characterized when a star-operation $\star$ is given by the map $A \mapsto A^\star = \cap AD_{P_\alpha}$, with $D = \cap D_{P_\alpha}$.

I am presently working on characterizing when a star-operation $\star$ is given by the map $A \mapsto A^\star = \cap AD_\alpha$, with $D = \cap D_\alpha$.

REFERENCES

- [1] D. D. Anderson, *Star-operations induced by overrings*, Comm. Algebra 16(1988), 2535-2553.
- [2] D. D. Anderson and D. F. Anderson, *Examples of star-operations on an integral domain*, Comm. Algebra 18(1990), 1621-1643.
- [3] D. D. Anderson and S. J. Cook, *Two star-operations and their induced lattices*, Comm. Algebra 28(2000), 2461-2475.
- [4] J. T. Arnold and J. W. Brewer, *On Flat Overrings, Ideal Transforms and Generalized Transforms of a Commutative Ring*, Journal of Algebra 18(1971), 254-263.
- [5] R. Gilmer, *Multiplicative Ideal Theory*, Queen's in Pure and Applied Mathematics, vol. 90, Queen's University, Kingston, Ontario, Canada, 1992.
- [6] F. Halter-Koch, *Ideal Systems, An Introduction to Multiplicative Ideal Theory*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 211, Marcel Dekker, Inc, New York, New York, 1998.
- [7] P. Jaffard, *Les Systèmes d'Ideaux*, Travaux et Recherches Mathématiques, vol. IV, Dunod, Paris, 1960.
- [8] M. D. Larsen and P. J. McCarthy, *Multiplicative Theory of Ideals*, Pure and Applied Mathematics, vol. 43, Academic Press, Inc, New York and London, 1971.

Sharon M. Clarke
Department of Mathematics
14 MacLean Hall
The University of Iowa
Iowa City, Iowa, 52242
E-mail: smclarke@math.uiowa.edu